

Some remarks on the Wu-Sprung potential. Preliminary report

Diego Dominici *

Department of Mathematics

State University of New York at New Paltz

75 S. Manheim Blvd. Suite 9

New Paltz, NY 12561-2443

USA

Phone: (845) 257-2607

Fax: (845) 257-3571

1 Introduction

Let $V(x)$ be the solution of the equation

$$\pi x = \sqrt{V - V_0} \ln \left(\frac{V_0}{2\pi e^2} \right) + \sqrt{V} \ln \left[\frac{\sqrt{V} + \sqrt{V - V_0}}{\sqrt{V} - \sqrt{V - V_0}} \right], \quad (1)$$

where $V_0 > 0$. Rationalizing the denominator, we can write (1) in the more convenient form

$$\pi x = \sqrt{V - V_0} \ln \left(\frac{V_0}{2\pi e^2} \right) + \sqrt{V} \ln \left[\frac{(\sqrt{V} + \sqrt{V - V_0})^2}{V_0} \right]. \quad (2)$$

Setting

$$V = V_0 \sec^2(\theta), \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2} \quad (3)$$

*e-mail: dominicd@newpaltz.edu

in (2), we obtain

$$\frac{\pi x}{\sqrt{V_0}} = \tan(\theta) \ln\left(\frac{V_0}{2\pi e^2}\right) + 2 \sec(\theta) \ln[\sec(\theta) + \tan(\theta)]. \quad (4)$$

It follows from (4) that $x = 0$ if $\theta = 0$ or $\theta = \theta_0$, with

$$V_0 = 2\pi e^2 [\sec(\theta_0) + \tan(\theta_0)]^{-2\sec(\theta_0)}. \quad (5)$$

In the first case $x = 0$ and $V = V_0$, and in the second $x = 0$ and $V = V_0 \sec^2(\theta_0) > V_0$.

Equation (5) admits a solution only when $0 < V_0 < 2\pi$ (see Figure 1). Thus, (2) generates a family of curves for $0 < V_0 < 2\pi$ and it defines a single valued function $V(x)$ for $V_0 \geq 2\pi$ (see Figure 2). This was observed before in [2], but without any justification.

2 Behavior for small x

In this section we determine the expansion of $V(x)$ in a neighborhood of the point $(0, V_0)$. We shall distinguish two cases, depending on $V_0 \neq 2\pi$ and $V_0 = 2\pi$.

2.1 Case I: $V_0 \neq 2\pi$

We consider the following expansion for $V(x)$

$$V(x) = V_0 + \sum_{k=1}^{\infty} a_k (\pi x)^{2k} \omega^{2k-1} (-V_0)^{1-k}, \quad (6)$$

where

$$\omega = \left[\ln\left(\frac{V_0}{2\pi}\right) \right]^{-1}. \quad (7)$$

Replacing (6) in (2) and expanding in powers of x , we obtain

$$\begin{aligned} a_1 &= \omega \\ a_2 &= \frac{4}{3}\omega^2 \end{aligned}$$




Figure 1: A sketch of V_0 as a function of θ_0 .




Figure 2: A sketch of $V(x)$ for various values of V_0 .

$$a_3 = \frac{8}{15}\omega^2 + \frac{28}{9}\omega^3$$

$$a_4 = \frac{32}{105}\omega^2 + \frac{16}{5}\omega^3 + \frac{80}{9}\omega^4$$

$$a_5 = \frac{64}{315}\omega^2 + \frac{528}{175}\omega^3 + \frac{704}{45}\omega^4 + \frac{2288}{81}\omega^5$$

$$a_6 = \frac{512}{3465}\omega^2 + \frac{13312}{4725}\omega^3 + \frac{14144}{675}\omega^4 + \frac{5824}{81}\omega^5$$

$$+ \frac{23296}{243}\omega^6$$

$$a_7 = \frac{1024}{9009}\omega^2 + \frac{12800}{4851}\omega^3 + \frac{13312}{525}\omega^4 + \frac{356864}{2835}\omega^5$$

$$+ \frac{8704}{27}\omega^6 + \frac{82688}{243}\omega^7$$

$$a_8 = \frac{4096}{45045}\omega^2 + \frac{17408}{7007}\omega^3 + \frac{5892608}{202125}\omega^4 + \frac{1901824}{10125}\omega^5$$

$$+ \frac{661504}{945}\omega^6 + \frac{578816}{405}\omega^7 + \frac{909568}{729}\omega^8$$

$$a_9 = \frac{8192}{109395}\omega^2 + \frac{136192}{57915}\omega^3 + \frac{153819136}{4729725}\omega^4 + \frac{599911168}{2338875}\omega^5$$

$$+ \frac{22685696}{18225}\omega^6 + \frac{13536512}{3645}\omega^7 + \frac{68913152}{10935}\omega^8 + \frac{30764800}{6561}\omega^9$$

$$a_{10} = \frac{131072}{2078505}\omega^2 + \frac{3670016}{1640925}\omega^3 + \frac{229421056}{6449625}\omega^4 + \frac{4575588352}{13820625}\omega^5$$

$$+ \frac{2095044608}{1063125}\omega^6 + \frac{1036288}{135}\omega^7 + \frac{208812032}{10935}\omega^8 + \frac{6735872}{243}\omega^9$$

$$+ \frac{117877760}{6561}\omega^{10}$$

and, in general, a_n is a polynomial of degree n in ω .

In Figures 3 and 4, we graph $V(x)$ and the first 10 terms of (6) with $V_0 = 3.1$ and $V_0 = 7.1$, respectively.




Figure 3: A comparison of $V(x)$ (solid curve) and its approximation, with $V_0 = 3.1$ (ooo).




Figure 4: A comparison of $V(x)$ (solid curve) and its approximation, with $V_0 = 7.1$ (ooo).

2.2 Case II: $V_0 = 2\pi$

Since $\omega = \infty$ when $V_0 = 2\pi$, the expansion (6) is no longer valid. Instead, we shall use

$$V(x) = 2\pi + \pi \sum_{k=1}^{\infty} b_k (3x)^{\frac{2k}{3}} 5^{1-k} \pi^{\frac{k}{3}}. \quad (8)$$

Replacing (8) in (2) and expanding in powers of x , we get

$$\begin{aligned} b_1 &= 1, & b_2 &= \frac{2}{3}, & b_3 &= \frac{1}{21}, & b_4 &= -\frac{2}{567}, & b_5 &= \frac{92}{130977} \\ b_6 &= -\frac{4}{21021}, & b_7 &= \frac{19543}{321810489}, & b_8 &= -\frac{352610}{16412334939} \\ b_9 &= \frac{12799}{1568439873}, & b_{10} &= \frac{6350075192}{1944910927276317}, \dots \end{aligned}$$

In Figure 5 we sketch $V(x)$ and the the first 10 terms of (8).

3 Asymptotic behavior

We shall now analyze the behavior of $V(x)$ as $x \rightarrow \infty$. From (2) we have

$$\sqrt{V - V_0} \ln \left(\frac{V_0}{2\pi e^2} \right) + \sqrt{V} \ln \left[\frac{(\sqrt{V} + \sqrt{V - V_0})^2}{V_0} \right] \sim \ln \left(\frac{2V}{\pi e^2} \right) \sqrt{V}$$

as $V \rightarrow \infty$. Therefore,

$$\ln \left(\frac{2V}{\pi e^2} \right) \sqrt{V} \sim \pi x, \quad x \rightarrow \infty,$$

which implies

$$V(x) \sim \frac{\pi^2 x^2}{4} \left[\text{LW} \left(\sqrt{\frac{\pi}{2}} \frac{|x|}{e} \right) \right]^{-2}, \quad |x| \rightarrow \infty \quad (9)$$

where $\text{LW}(\cdot)$ represents the Lambert-W function [1]. Note that the asymptotic behavior is independent of V_0 , as suggested by Figure 2.

In Figure 6 we graph $V(x)$ and its asymptotic approximation (9).




Figure 5: A comparison of $V(x)$ (solid curve) and its approximation, with $V_0 = 2\pi$ (ooo).




Figure 6: A comparison of $V(x)$ (solid curve) and its asymptotic approximation (ooo).

References

- [1] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth. On the Lambert W function. *Adv. Comput. Math.*, 5(4):329–359, 1996.
- [2] H. Wu and D. W. L. Sprung. Riemann zeros and a fractal potential. *Phys. Rev. E (3)*, 48(4):2595–2598, 1993.